

FUSE OBSERVATIONS OF THE ACTIVE COOL STAR AB DORADUS

T. B. AKE

Department of Physics & Astronomy, The Johns Hopkins University, and Computer Sciences Corporation,
3400 N. Charles Street, Baltimore MD, 21218

A. K. DUPREE AND P. R. YOUNG

Harvard-Smithsonian Center for Astrophysics, Cambridge MA 02138

J. L. LINSKY

JILA, University of Colorado and NIST, Boulder CO 80309-0440

R. F. MALINA, N. W. GRIFFITHS, AND O. H. W. SIEGMUND

Space Sciences Laboratory, University of California, Berkeley, Berkeley CA 94720

AND

B. E. WOODGATE

Laboratory for Astronomy and Solar Physics, NASA's GSFC, Code 681, Greenbelt MD 20771

1 June 2000

ABSTRACT

Far ultraviolet spectra were obtained of the active cool star AB Doradus (HD 36705) during the calibration and checkout period of the *FUSE* satellite. Observations in this early phase of the mission were taken at a resolving power of 12000–15000 (~ 20 – 25 km s^{−1}) and covered the spectral range 905–1187 Å. The integrated spectrum exhibits strong, rotationally broadened stellar emission from C III ($\lambda 977$, $\lambda 1175$) and O VI ($\lambda 1032$, $\lambda 1037$), and many weaker lines. Strong emission lines of C III and O VI exhibit broad wings. The C III $\lambda 977$ profile shows blue-shifted absorption at ~ 30 km s^{−1} and C II $\lambda 1036$ absorption appears superposed on emission in the wing of O VI $\lambda 1037$. Rotational modulation of C III and O VI is present, in harmony with its photometric variability. Flares were detected in the brightest lines and subexposures were analyzed to examine flux and profile variations. Downflows that extend to 600 km s^{−1} during a flare are found in the O VI profiles. These early observations demonstrate that *FUSE* will be an exceptional instrument for studying chromospheres in cool stars.

Subject headings: stars:chromospheres – stars:flare – stars:individual(AB Dor) – stars:pre-main-sequence – ultraviolet:stars

1. OBSERVATIONS AND DATA ANALYSIS

AB Doradus is a nearby (15 pc), rapidly rotating ($P_{\text{rot}} = 12.4$ hr, $v \sin i = 90$ km s^{−1}), K0–2 IV–V pre-main sequence star that exhibits high activity and frequent flaring (e.g., Collier Cameron et al. 1999; Schmitt, Cutispoto & Krautter 1998; Vilhu et al. 1998). It is the first cool star observed by the *Far Ultraviolet Spectroscopic Explorer* (*FUSE*, cf. Moos et al. 2000). Observations were obtained on 1999 October 20 and 22, in the LiF channels ($\lambda\lambda 987$ – 1187), and on 1999 December 14, when exposures in all four channels were made covering the range $\lambda\lambda 905$ – 1187 . Exposures were taken in time-tagged mode at 1 second resolution through the $30'' \times 30''$ aperture and were scheduled when AB Dor was in the continuous viewing zone. Thus good coverage was obtained over most of its rotational period each day. The only interruptions were due to South Atlantic Anomaly crossings, when detector high voltage is reduced. The total exposure time amounted to 79 ks for the LiF channels and 24 ks for SiC, of which 33 ks and 13 ks respectively were obtained during orbital night. The spectral resolving power was 12000–15000 (~ 20 – 25 km s^{−1}) but the considerable oversampling improves the spectral purity.

Since the spectrographs were not fully focused, care was exercised when comparing lines from different channels. Data sets were reprocessed with the calibration pipeline as improvements became available. Both airglow and the

scattered solar spectrum can contribute emission lines expected to be present in cool stars; thus spectra obtained from data taken during orbital night were examined and analyzed separately in regions where the contamination was significant. For time variability studies, subexposures were created from the photon lists and reprocessed individually.

Wavelength calibration errors especially near the ends of detector segments are still present. For AB Dor, the easily identifiable lines and airglow lines allowed for local adjustments of the scale, but identifications for weaker lines had to be confirmed through comparisons with solar disk spectra from SUMER (e.g., Curdt et al. 1997). The wavelength scale uncertainty precludes at present an analysis of possible differential motions between the lines.

2. LINE SPECTRUM

The *FUSE* spectrum of AB Dor is shown in Figure 1. The strongest lines arise from C III ($\lambda 977$, $\lambda 1175$), O VI ($\lambda 1032$, $\lambda 1037$) and airglow lines of H I, O I, N I, and N II. Most of the latter can be distinguished from stellar components by screening out parts of the exposures taken in daylight as shown in Fig. 1. We can tentatively identify fainter lines due to H I, He II, C I, C II, N III, Si III, Si IV, S III, S VI, and Fe III, which will be discussed in a later paper. The lines are broadened by the high rotation rate of the star.




FIG. 1.— *FUSE* spectra (SiC1A, SiC1B, SiC2B, LiF1B, LiF2B) of AB Dor taken on 1999 December 14 with prominent stellar and airglow emission features identified. The upper spectrum in each panel includes photons collected during orbital day and night; the lower spectrum is derived from night time photons only. Rotational broadening is apparent even in these compressed plots.

Profiles of chromospheric and transition region lines can provide clues to heating mechanisms and mass motions in cool stars. With a resolving power of 12000–15000, *FUSE* can be used to study thermal and non-thermal motions, which typically range from 15 to hundreds of km s^{-1} in these stars (e.g., Wood, Linsky, & Ayres 1997). Figure 2 shows the O VI doublet region in detail for the time-integrated LiF1A spectrum on October 22. The $\lambda 1032$ component is cleanly separated from other features in the spectrum and is symmetrical. A Gaussian profile broadened by 90 km s^{-1} rotation with no limb darkening can adequately reproduce the core of the line, but does not fit the wings. Schmitt et al. (1997) were able to measure the rotational broadening in O VI with *ORFEUS*, but could not adequately discern the broad wings. The low scattered light properties of *FUSE* (Sahnou et al. 2000) and the relatively modest FUV count rates for AB Dor make an instrumental source unlikely to explain the extended profile. Extra flux in the wings of Si IV and C IV lines has been observed in other active stars (Wood, Linsky, & Ayres 1997), and the O VI observations of AB Dor extend this phenomenon to greater line-formation temperatures. Since telescope focus and spectrograph calibration had not been completed, there is some disagreement in the details of the line profiles between the *FUSE* channels. Considering all the data, we find a two-Gaussian fit

composed of a narrow ($\text{FWHM} \simeq 80 \text{ km s}^{-1}$) and broad ($\text{FWHM} \simeq 300 \text{ km s}^{-1}$) component, the latter contributing 38–56% of the total line flux depending on the data set, reproduces the $\lambda 1032$ profile when rotationally and instrumentally broadened. These values are comparable to widths of 68 and 334 km s^{-1} determined for C IV in AB Dor from GHRs spectra (Vilhu et al. 1998). Wood, Linsky, & Ayres (1997) found broad components in a number of cool stars to be correlated with C IV and X-ray surface fluxes, and proposed that their origin is microflare heating. However extended material may occur frequently in rapidly rotating cool stars. Co-rotating prominence material or an extended chromosphere as Collier Cameron & Robinson (1989) have identified on AB Dor might cause similar broadening. Another rapidly rotating K dwarf, V471 Tau, exhibits material at 10^5 K extending to $1 R_*$ above the surface (Guinan et al. 1986).

The weaker member of the O VI doublet ($\lambda 1037.61$) is blended with C II emission ($\lambda 1036.34$, $\lambda 1037.02$) in its blue wing. A strong C II $\lambda 1036$ absorption feature is superposed on the emission, likely due to interstellar or circumstellar material. Except for the blue wing, the O VI $\lambda 1037$ line is well matched by the same Gaussian fit from $\lambda 1032$ when shifted by the difference in wavelengths and scaled by the doublet ratio.

C III $\lambda 977$ differs from O VI by additional absorption ap-

pearing $\sim 30 \text{ km s}^{-1}$ from line center and irregularities in the wings of the line profile (Figure 3). We have considered several possible origins for this, which cannot be resolved until a better wavelength scale is available. A similar asymmetric profile in C III was found in the luminous cool star β Dra from *ORFEUS* spectra where it was attributed to a wind (Dupree & Brickhouse 1998). AB Dor frequently displays transient absorption features in the H α profile (Collier Cameron et al. 1999), and the C III absorption may be a high temperature manifestation of the behavior in H α . Finally, the presence of the C II $\lambda 1036$ absorption line suggests that the C III absorption could also be interstellar or circumstellar in origin.

3. ELECTRON DENSITIES

We compute a ratio of the density sensitive C III lines of, $\lambda 1175/\lambda 977 = 0.651 \pm 0.172$ (in erg units), although the absolute and relative flux calibration of *FUSE* is still at an early stage. This ratio, when compared to the calculations of the CHIANTI code, suggest that the density at $T \sim 80000 \text{ K}$ is near the high density limit, namely $\sim 10^{11} \text{ cm}^{-3}$, and similar to the values derived from *ORFEUS* spectra of AB Dor (Schmitt, Cutispoto, & Krautter 1998). However the uncertain calibration and presence of absorption in the $\lambda 977$ line, potentially weakening the line and increasing the inferred density, suggests that caution must be used in applying this diagnostic.

4. UV FLARES ON AB DOR

AB Dor undergoes frequent flaring, particularly in X rays; impulsive flares have also been detected in C IV (Vilhu et al. 1998). Several flares occurred during the *FUSE* observations; one was bright enough on 1999 October 22 to extract a spectrum in the strongest lines. Figure 4 shows a dynamic spectrum in the C III $\lambda 1175$ region. Light curves for C III $\lambda 1175$ and O VI show similar temporal behavior marking a short-duration ($\approx 700 \text{ sec}$) flaring episode. The overall brightening of the $\lambda 1175$ line was accompanied by an enhanced red shifted component. Here care was taken in determining which features are stellar flares as distinguished from events bursts seen with the detector (Sahnou et al. 2000). Fig. 2 shows the spectrum of red shifted components in the O VI lines; the Doppler shift of a Gaussian fit to the flare component is centered at $+235 \text{ km s}^{-1}$, with extension to $+600 \text{ km s}^{-1}$. A flare in C IV observed by Vilhu et al. (1998) exhibited a broad blue-shifted Gaussian profile superposed on the quiescent profile and was attributed to evaporation of chromospheric material. Flares on AD Leo have shown high redshifts in C IV, amounting to $\sim 650 \text{ km s}^{-1}$ with extensions to 1800 km s^{-1} (Bookbinder et al. 1992). Downward flowing flare material is a frequent occurrence in solar post-flare magnetic loops and could explain the profiles observed here.

5. ROTATIONAL MODULATION

AB Dor undergoes optical and UV variations as active structures rotate through the line of sight. Maximum spottedness occurs at minimum light in V and U. In Figure 5, we show the variation of the strongest lines (C III and

O VI) on 1999 October 22, when AB Dor appears to have been most active, by dividing the *FUSE* observations into subexposures of $\sim 35 \text{ min}$ apiece. Rotational phase is determined from the ephemeris of Innis et al. (1988): $\text{HJD} = 2444296.575 + 0.51479 \text{ E}$. In this case, minimum light occurs at $\phi = 0.5$. Photometry in 1999 December from Mt. Kent Observatory in Queensland confirms that AB Dor continues to be faint at $\phi = 0.5$ on this ephemeris, although coverage over the entire rotational period was not obtained.

A peak in the *FUSE* fluxes occurs near $\phi \sim 0.6$; this would be expected from enhanced high temperature activity due to star spots on the photosphere and overlying regions of activity. Comparable modulation amplitudes in C III and O VI suggest the spatial distribution of the corona is similar between 80000 K and $2 \times 10^5 \text{ K}$ where these lines are formed. Greater enhancement of C III than O VI during the flare may be affected by the increased density enhancing the metastable level of C III. The large flare on this date, however, occurs after the photometric minimum, near $\phi = 0.9$ suggesting that it arose from an active region not associated with the maximal spot coverage.

Simultaneous *ORFEUS* and *ROSAT* observations are discussed by Schmitt, Cutispoto, & Krautter (1998), who found that FUV and X-ray variations are correlated but do not appear to be rotationally modulated, except at about the 15% level for X-ray emission. AB Dor is known to undergo night-to-night optical variations (Collier Cameron et al. 1999), so O VI rotational modulation could be a transient phenomenon dependent on the activity of the star. For the X-ray emission, either the coronal material is more homogeneously distributed than the lower temperature material or the activity is at higher latitudes, and hence, always visible. We note that Collier Cameron et al. (1999) find AB Dor has a stable polar cap of photospheric spots.

6. CONCLUSIONS

This first set of observations of a cool star demonstrates that *FUSE* can provide insight into a variety of phenomena on AB Dor. We have identified lines spanning more than a decade in line formation temperature that can be used to determine diagnostics in the far ultraviolet. Even though the instrument is not fully focused, the line profiles can be studied in detail, and we have begun examining the dynamics and heating processes in this star. Through time-resolved spectroscopy, we can study variability on different time scales, from seconds to days. Importantly, the redundant wavelength coverage from different channels provides high confidence in observed variations of features and structures.

This work is based on data obtained for the Guaranteed Time Team by the NASA-CNES-CSA FUSE mission operated by The Johns Hopkins University. Financial support to U.S. participants has been provided by NASA Contract NAS5-32985. Photometric observations during 1999 December were kindly provided by Ian Waite and Brad Carter of the University of South Queensland, Australia.

REFERENCES

- Bookbinder, J., Walter, F., & Brown, A. 1992 in ASP Conf. Ser., 26, Cambridge Workshop on Cool Stars, Stellar Systems and the Sun, ed. M. S. Giampapa & J. A. Bookbinder (San Francisco:ASP), 27
- Collier Cameron, A., & Robinson, R. D. 1989, MNRAS, 236, 57
- Collier Cameron, A. et al. 1999, MNRAS, 308 493
- Curdt, W., Feldman, U., Laming, J. M., Wilhelm, K., Schühle, U., & Lemaire, P. 1997, A&AS, 126, 281
- Dupree, A. K., & Brickhouse, N. S. 1998, ApJ, 500, L33
- Guinan, E. F., Wacker, S. W., Baliunas, S. L., Loeser, J. G., & Raymond, J. C. 1986, ESA SP-263, p. 197
- Innis, J. L., Thompson, K., Coates, D. W., & Lloyd Evans, T. 1988, MNRAS, 235, 1411
- Moos, H. W., et al. 2000, ApJ, this volume
- Sahnow, D. J., et al. 2000, ApJ, this volume
- Schmitt, J. H. M. M., Cutispoto, G., & Krautter, J. 1998, ApJ, 500, L25
- Schmitt, J. H. M. M., et al. 1997, A&A, 325, 249
- Vilhu, O., Muhli, P., Huovelin, J., Hakala, P., Rucinski, S. M., & Collier Cameron, A. 1998, AJ, 115, 1610
- Wood, B. E., Linsky, J. L., & Ayres, T. R. 1997, ApJ, 478, 745




FIG. 2.— *Upper panel:* The O VI doublet, time-integrated over the 1999 October 22 exposures with the LiF1A channel. A two-component Gaussian fit is required for the $\lambda 1032$ line. Shown are the Gaussians after rotational and instrumental broadening (dotted lines), the final fit (thick line), and the same profile shifted and scaled by a factor of two for $\lambda 1037$ (dashed line). Stellar C II emission is noted shortward of O VI $\lambda 1037$, with interstellar or circumstellar C II $\lambda 1036$ absorption superimposed. *Lower panel:* Flare profiles are shown averaged over a 2500 s active period of the star. Red shifted material at $+235$ km s⁻¹, extending to 600 km s⁻¹, is found in O VI, as well as sharp emission components near line center. Dashed line is the two-component fit from the time-integrated profile in the upper panel.




FIG. 3.— Comparison of profiles of the O VI and C III lines showing the short wavelength absorption feature in C III, and broadened “bumpy” wings of emission in the C III profile. The velocity scale has an arbitrary offset for each line, and was simply coaligned in this figure.




FIG. 4.— Dynamic spectra of C III $\lambda 1175$ on 1999 October 22 totaled in 10 second bins. Small precursor flares occurred before the bright event at 05:28UT where both increased flux and red shifted emission are found. Gaps in data are due to SAA passages. The feature showing diurnal variation at $\lambda 1168$ is second-order scattered solar He I.




FIG. 5.— Variation of the O VI $\lambda 1032$, $\lambda 1037$ and C III $\lambda 1175$ line fluxes with rotational phase for 1999 October 22 observations binned in 35 minute intervals. The measurements are replicated over two rotational phases. Spot maximum is expected at $\phi \sim 0.5$. The large flare at 05:28 UT occurred at $\phi = 0.9$.